

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A23K 1/16, 1/18		A1	(11) International Publication Number: WO 00/51443 (43) International Publication Date: 8 September 2000 (08.09.00)
(21) International Application Number: PCT/US00/05586 (22) International Filing Date: 3 March 2000 (03.03.00) (30) Priority Data: 60/122983 5 March 1999 (05.03.99) US 09/518116 3 March 2000 (03.03.00) US (71) Applicant: THE IAMS COMPANY [US/US]; 7250 Poe Avenue, Dayton, OH 45414 (US). (72) Inventors: DAVENPORT, Gary, Mitchell; 4020 Shell Avenue, Dayton, OH 45415 (US). HAYEK, Michael, Griffin; 926 Tammy Circle, Dayton, OH 45415 (US). (74) Agents: LUNA, Susan, M. et al.; Killworth, Gottman, Hagan & Schaeff, L.L.P., One Dayton Centre, Suite 500, One South Main Street, Dayton, OH 45402-2023 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: PROCESS FOR PRESERVING SKELETAL MUSCLE MASS IN GERIATRIC DOGS			
(57) Abstract A process is provided for preserving skeletal muscle mass in a geriatric dog by feeding the dog a diet containing an effective amount of animal-based protein. Preferably, the diet includes greater than about 16 % by weight animal-based protein on a dry matter basis, and more preferably, from about 24 to 34 % by weight protein. Such a diet also provides sufficient protein to improve the body composition of the dog and reduce body fat.			

858

Atkin

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PROCESS FOR PRESERVING SKELETAL MUSCLE MASS IN GERIATRIC DOGS

This invention relates to a method for preserving skeletal muscle mass in geriatric dogs, and more particularly, to such a method which includes providing a diet which includes beneficial amounts of animal-based protein to preserve skeletal muscle mass, reduce body fat and improve the overall body composition of geriatric dogs.

Advanced age in mammals is often accompanied by a loss of skeletal muscle mass and a loss of muscle strength. Muscle growth is the result of protein synthesis and protein degradation. Therefore, decline in muscle mass may be due to an imbalance in the protein synthesis/protein degradation ratio of skeletal muscle protein.

Various muscle-specific enzymatic processes may play a role in regulating protein turnover and protein degradation in skeletal muscle such as the ATP-dependent ubiquitin-proteasome pathway and the calcium-dependent enzyme μ -calpain and its endogenous inhibitor calpastatin.

Other factors which may affect loss of skeletal muscle mass and strength include age-related changes in the contractile mechanism. A major regulatory component of the contractile mechanism is troponin-T, which is one of three subunits of the troponin protein molecule which includes troponin-I and troponin-C. Other muscle-specific proteins include actin, myosin, and tropomyosin. Troponin and tropomyosin control the myosin-actin interactions involved in calcium-mediated muscle contractions. Troponin-I inhibits the interaction of actin and myosin in non-contracted muscle, while troponin-C contains a calcium-binding site that initiates muscle contractions. Troponin-T functions to bind the other troponin subunits to tropomyosin.

Variations in troponin-T isoform expression may be related to differences in calcium sensitivity of tension development and maximal velocity of shortening of skeletal muscle fibers. Therefore, variations in troponin-T isoform expression may explain some loss of muscle strength and tone. Differences in myofibrillar protein isoform expression may also impact the rate of protein turnover as different isoforms may exhibit variable susceptibilities to proteolysis.

Accordingly, there is a need in the art for a method of regulating skeletal muscle protein in animals such as geriatric dogs to reduce muscle protein degradation and the loss of skeletal muscle mass.

The present invention addresses that need by providing a process for
5 preserving skeletal muscle mass in geriatric dogs by feeding the dogs a diet containing an effective amount of animal-based protein. By geriatric dogs, it is meant those dogs which have reached a senior age associated with their breed and/or have begun to exhibit signs of aging. For giant breeds, the senior stage is considered to begin between about 5 to 7 years of age; for medium breeds, about 7 to 9 years; and for
10 small breeds, between about 8 to 10 years.

Preferably, the process comprises feeding a geriatric dog a diet including greater than about 16% by weight protein on a dry matter basis, and more preferably, from about 24 to 34% by weight, where the protein is provided in the diet as an animal-based protein source.

15 Preferably, the diet comprises, on a dry matter basis, about 24 to 34% by weight protein, about 9 to 22% by weight fat, and about 1 to 6% by weight dietary fiber.

The diet of the present invention has also been found to reduce body fat in geriatric dogs and to improve the body composition of geriatric dogs. By body composition, it is meant the total quantity of lean, fat and bone in the body. By
20 improved body composition, it is meant that the dogs exhibit a greater percentage of lean tissue, and a lower percentage of body fat.

Accordingly, it is a feature of the invention to provide a process for preserving skeletal muscle mass in geriatric dogs by providing an effective amount of animal-based protein in the diet of the animals. It is a further feature of the invention to
25 provide a process for improving the body composition and reducing the body fat of geriatric dogs. These, and other features and advantages of the present invention, will become apparent from the following detailed description and accompanying drawings.

Fig. 1 is an immunoblot showing myofibrillar protein fraction from geriatric and young canines fed diet A;

Fig. 2 is an immunoblot showing myofibrillar protein fraction from geriatric and young canines fed diet B; and

5 Fig. 3 is an immunoblot showing myofibrillar protein fraction from geriatric and young canines fed diet C.

The present invention provides a dietary composition which contains a source of animal-based protein in an amount greater than about 16% protein by weight on a dry matter basis, and preferably between about 24 to 34% by weight. Such a diet has
10 been found to produce a shift in the troponin-T isoform expression in old dogs which is similar to the troponin-T expression of younger dogs. It is believed that the dietary composition of the present invention, when fed to geriatric dogs, results in a reduction in skeletal muscle protein degradation and a preservation of skeletal muscle mass.

The dietary composition may be provided in the form of any suitable pet food
15 composition which also provides adequate nutrition for the animal. For example, a preferred canine diet for use in the present invention contains from about 24 to 34% by weight protein, 9 to 22% by weight fat, and about 1 to 6% by weight dietary fiber.

Preferably, chicken is the primary dietary protein source; however, other suitable animal protein sources include lamb, beef, fish, duck, deer, rabbit, and pork.

20 In order that the invention may be more readily understood, reference is made to the following examples which are intended to illustrate the invention, but not limit the scope thereof.

Example 1

25 Twenty-three female beagles averaging 4 years of age or less (n=12) and over 9 years of age (one 11 year old, one 12 year old and ten 10 year old) (n=12) were fed diets containing either 16% (Diet A) or one of two 32% protein formulations (four dogs per age group per diet). The two 32% protein formulations differed in the amount of animal-based protein. The low animal-based protein diet (Diet B) used chicken protein at the same level as was used for the 16% diet with the balance of the protein from
30 plant protein. The high animal-based protein diet (Diet C) used chicken as the primary dietary protein source. The composition of diets A, B and C is shown below in Table 1.

TABLE 1
Diet Composition of Experimental Diets

Ingredient	Diet A (16% CP, meat)	Diet B (32% CP, meat +veg.)	Diet C (32% CP, meat)
5 Refined poultry meal	18.48	18.99	39.98
Corn gluten meal	0	23.94	0
Ground corn	16.49	33.43	33.58
Corn starch	37.85	0	5.75
Poultry fat	11.34	9.45	8
10 Poultry digest	3	3	3
Beet pulp	4	4	4
Dicalcium phosphate	3.2	2.55	1.15
Menhaden oil	1.1	1.1	1.1
15 Brewer's yeast	1	1	1
Vitamin premix	0.9	0.35	0.35
Potassium chloride	0.75	0.45	0.25
Calcium carbonate	0.65	0.75	0.55
Ground flax	0.3	0.3	0.3
20 Choline chloride	0.3	0.1	0.15
Sodium chloride	0.25	0.25	0.25
DL-methionine	0.2	0.15	0.4
Mineral premix	0.2	0.2	0.2

Nutrient analysis %	Diet A	Diet B	Diet C
Protein	15.41	31.35	33.02
Fat	16.43	16.82	17.56
NFE (starch)	65	66.66	66.99
Moisture	9.77	8.02	5.97
Ash	6.2	6.36	7.4
Fiber	2.58	2.14	2.08
Gross energy (kcal/gm)	4514.08	5010.01	4947.08

The dogs were limit fed to maintain metabolic body size. Prior to the start of the experimental diets, the dogs were fed a 24% animal-based protein diet for approximately 60 days. Immediately preceding the time that the dogs were switched to the experimental diets, a muscle biopsy sample was taken. A second muscle biopsy was taken after 45 days on the experimental diets. Three animals were excluded from the young age group. Animal 1 was excluded because it was 4 years of age, making it significantly older than the other animals (all other young dogs were approximately 1 year of age). Animals 2 and 3 were excluded from the current study because of sampling error. This left three young dogs on diet A, four young dogs on diet B and three young dogs on diet C. There still remained four geriatric dogs on each of the three dietary treatments.

Semitendinosus muscle biopsy samples were immediately frozen in liquid nitrogen after being removed from anesthetized dogs by a licensed veterinarian. These samples were then stored at -80°C until they were used for analysis.

In order to determine the state of the calpastatin molecules, the sarcoplasmic protein fraction from the biopsy samples was isolated and used for chemiluminescent immunoblotting analyses using commercially available monoclonal antibodies against calpastatin. Briefly, a 0.1 gram sample was homogenized with a motor driven Dounce homogenizer in five volumes of extraction buffer (10 mM EDTA, 100 mg/L trypsin inhibitor, 2 µM E-64, 3 mM PMSF and 0.1% B-mercaptoethanol, 100 mM Tris, pH 8.5). Samples were homogenized over three intervals of 20 strokes each for 30 seconds. Homogenized samples were removed from the homogenizers and were transferred

into microcentrifuge tubes by using a transfer pipette and rinsing the homogenizing tubes with 100 μ l of extraction buffer. Samples were centrifuged for 20 minutes at 10,850 x g at 4°C. After centrifugation, the supernatant was removed from the pellet using a 1 cc glass syringe with a 22 gauge needle. The supernatant was then dialyzed
5 overnight at 8°C against a solution containing 1 mM EDTA, 0.1% B-mercaptoethanol and 40 mM Tris pH 7.5. After dialysis, the protein concentration of the supernatant was determined using the Bradford assay.

Samples for gel electrophoresis were prepared as follows. One hundred micrograms of protein were loaded into each well of a 15% acrylamide separating gel
10 (15% acrylamide/bis, [100:1 acrylamide:bisacrylamide], .375 M Tris-HCl, pH 8.8, 0.1% sodium dodecyl sulfate (SDS), 0.1% ammonium persulfate and 0.67% N,N,N',N'-Tetramethylethylenediamine (TEMED) with a 5% stacking gel (4% acrylamide/bis, [100:1 acrylamide:bisacrylamide], 0.125 M Tris-HCl, pH 6.8, 0.1% SDS, 0.125 TEMED and 0.75% ammonium persulfate. The composition of the running
15 buffer used was 25 mM Tris, 192 mM glycine and 0.1% SDS. Gels were run at 120 V for approximately 195 minutes (until the dye front had run off of the gel). Immediately after the run, the proteins on the gels were transferred to PVDF membranes that had been prewet in 100 % methanol and then in transfer buffer (25 mM Tris, 192 mM glycine and 15% methanol). Gels were transferred for 90 minutes at 90 volts. After
20 transfer, the membranes were blocked for one hour at room temperature in a 5% solution of non-fat dry milk dissolved in PBS-Tween (80 mM Na_2HPO_4 , 20 mM NaH_2PO_4 , 100 mM NaCl and 0.1% Tween). After blocking, the membranes were incubated in primary antibody (cat # RDI-CALPSTabm, mouse anti-calpastatin, Research Diagnostics Inc. Flanders, NJ) at a dilution of 1:2000 (PBS-Tween:antibody).
25 The primary incubation was done overnight at 8°C. After primary incubation, the membranes were allowed to reach room temperature and were then washed three times (ten minutes per wash) in PBS-Tween. After rinsing, the secondary antibody (Goat-anti-mouse IgG conjugated with horseradish peroxidase, product #A2554, Sigma Chemical Co., St. Louis Mo.) was applied at a dilution of 1:5000
30 (PBS-Tween:antibody). Incubation in secondary antibody was done at room temperature for one hour. Membranes were then washed three times (ten minutes per wash) in PBS-Tween. The presence of calpastatin was detected using ECL Western

Blotting reagents (Amersham Life Science, Arlington Heights, IL) as directed by the manufacturer.

The myofibrillar fraction from selected samples was also subjected to immunoblotting techniques to determine if there was a difference in degradation or in isoform expression of specific μ -calpain substrate proteins. The samples for gel electrophoresis were prepared according to the procedure outlined in Huff-Lonergan et al., "Proteolysis of Specific Muscle structural Proteins by μ -calpain at low pH and Temperature is Similar to Degradation in Postmortem Bovine Muscle", *J. Anim. Sci.* 74:993, 1996 and in Huff-Lonergan et al., "Sodium Dodecyl Sulfate Polyacrylamide Gel Electrophoresis and Western Blotting Comparisons of Purified Myofibrils and Whole Muscle Preparations for Evaluating Titin and Nebulin in Postmortem Bovine Muscle", *J. Anim. Sci.* 74:779, 1996. Gel electrophoresis, immunoblotting and detection of troponin-T were performed as described in Huff-Lonergan et al.

Results

Calpastatin

Careful examination of the immunoblots revealed that age had a significant effect upon the intensity of the band that corresponds to intact calpastatin, while dietary treatment did not. Older animals in this study exhibited a much greater intensity of the intact calpastatin band in samples taken both before and after the dietary treatment. This indicates that differences in level of expression and/or differences in modification of calpastatin play an important role in the physiological processes that accompany physiological aging.

Troponin-T

There are numerous isoforms of troponin-T that can be expressed in skeletal muscle. Within this experiment, the pattern of isoform expression of troponin-T was noted to differ in animals of different ages and across dietary treatments. When immunoblots probed for troponin-T were examined, it was apparent that age significantly impacted isoform expression in samples taken both before and after the dietary treatment (see Figures 1, 2, and 3; young vs. old). In samples that were taken

at the conclusion of the experimental dietary treatment, it was also noted that diet significantly impacted the expression of the troponin-T isoform pattern. In addition, there was a significant interaction between age and diet. When the immunoblots for the individual combinations of diet and treatment were examined, it was seen that the

5 geriatric canines that were fed diet C showed a shift in their isoform expression pattern. This shift was from that pattern that was expressed in nearly all geriatric canines prior to dietary treatment to an isoform pattern that more closely resembled that seen in the younger canines. This indicates that animal-based protein diets that exceed the level of protein in diet A affect age-induced changes in molecular and cellular events

10 associated with aging canine muscle.

Body Composition

The body composition of the dogs was assessed using dual-energy X-ray absorptiometry (DEXA) which estimates the total quantity of lean, fat and bone in the body.

TABLE 2

Old				Young			
	Diet A 16% protein	Diet B 32% protein	Diet C 32% protein	Diet A 16% protein	Diet B 32% protein	Diet C 32% protein	P<
Week 0							
% lean	73.76	79.22	76.24	81.37	82.80	84.53	
% fat	22.28	17.49	20.56	14.74	13.59	11.61	
% ash	3.96	3.30	3.21	3.89	3.62	3.86	
Week 6							
% lean	73.85 ^a	83.30 ^b	80.88 ^b	82.73 ^b	83.67 ^b	85.71 ^b	0.10
% fat	22.00 ^a	13.08 ^b	15.62 ^{ab}	13.59 ^b	12.53 ^b	10.34 ^b	0.10
% ash	4.16	3.63	3.50	3.68	3.80	3.94	
Absolute change (week 6 vs. week 0)							
% lean	.08 ^a	4.08 ^b	4.64 ^b	1.36 ^{ab}	.87 ^{ab}	1.18 ^{ab}	0.10
% fat	-.29 ^a	-4.41 ^b	-4.94 ^b	-1.15 ^{ab}	-1.06 ^{ab}	-1.26 ^{ab}	0.10
% ash	.20	.33	.30	-.22	.18	.08	

DEXA results showed that after 6 weeks of feeding, geriatric dogs fed diets B and C had greater ($P<0.05$) percentage of lean tissue and lower ($P<0.05$) percentage of body fat compared with dogs fed diet A. These changes in body composition were attributed to the effect of high protein diets on the preservation of lean body mass and loss of body fat. In contrast, no differences were noted after 6 weeks in the younger dogs.

Overall, the changes in skeletal muscle protein degradation in geriatric dogs associated with the high animal protein-based diets were reflected in improved body composition compared with dogs fed the lower protein diet or the protein diet which combined animal and vegetable protein sources.

While certain representative embodiments and details have been shown for purposes of illustrating the invention, it will be apparent to those skilled in the art that various changes in the methods and apparatus disclosed herein may be made without departing from the scope of the invention, which is defined in the appended claims.

CLAIMS

1. A process for preserving skeletal muscle mass in a geriatric dog comprising the step of feeding said dog a diet including greater than about 16% by weight animal-based protein on a dry matter basis.
- 5 2. The process claimed in claim 1 in which said diet includes from between about 24 to 34% by weight protein on a dry matter basis.
3. The process claimed in claim 1 in which said diet comprises, on a dry matter basis, about 24 to 34% by weight protein, about 9 to 22% by weight fat, and about 1 to 6% total dietary fiber.
- 10 4. A process for improving the body composition of a geriatric dog comprising the step of feeding said dog a diet including greater than about 16% by weight animal-based protein on a dry matter basis.
5. The process claimed in claim 4 in which said diet includes from between about 24 to 34% by weight protein on a dry matter basis.
- 15 6. A process for reducing body fat of a geriatric dog comprising the step of feeding said dog a diet including greater than about 16% by weight animal-based protein on a dry matter basis for a time sufficient to reduce body fat in said dog.
7. The process claimed in claim 6 in which said diet includes from between about 24 to 34% by weight protein on a dry matter basis.
- 20 8. The process claimed in claim 6 in which said diet comprises, on a dry matter basis, about 24 to 34% by weight protein, about 9 to 22% by weight fat, and about 1 to 6% total dietary fiber.

-11-

9. The process claimed in claim 1 wherein said animal-based protein is selected from the group consisting of chicken, lamb, beef, fish, duck, deer, rabbit, pork, and mixtures thereof.

10. The process claimed in claim 9 wherein said animal-based protein comprises
5 chicken.

Fig. 1/3

Diet A

Old

1 2 3 4 5 6 7 8



Young

1 2 3 4 5 6



Fig. 2/3
Diet B

Old

1 2 3 4 5 6 7 8



Young

1 2 3 4 5 6 7 8



Fig. 3/3
Diet C

Old

1 2 3 4 5 6 7 8



Young

1 2 3 4 5 6



INTERNATIONAL SEARCH REPORT

Int. national Application No

PCT/US 00/05586

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A23K1/16 A23K1/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A23K A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data, CHEM ABS Data, CAB Data, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	C.C. WILLIAMS ET AL.: "Effect of dietary protein on whole-body protein turnover and endocrine functionality in geriatric and young-adult dogs" FASEB JOURNAL., vol. 12, no. 5, 1998, page A878 XP002142344 FED. OF AMERICAN SOC. FOR EXPERIMENTAL BIOLOGY, BETHESDA, MD., US ISSN: 0892-6638 page A878, column 2, abstract 5085 --- -/--	1-10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"S" document member of the same patent family

Date of the actual completion of the international search

12 July 2000

Date of mailing of the international search report

08/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Dekeirel, M

INTERNATIONAL SEARCH REPORT

In. ational Application No

PCT/US 00/05586

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
P,X	E. HUFF-LONERGAN ET AL.: "Impact of age and diet on calpastatin and tropinin-T in young and old dogs" FASEB JOURNAL., vol. 13, no. 5, 1999, page A938 XP002142345 FED. OF AMERICAN SOC. FOR EXPERIMENTAL BIOLOGY, BETHESDA, MD., US ISSN: 0892-6638 page A938, column 1, abstract 690.11 ---	1-10
A	M.G. HAYEK ET AL.: "Nutrition and aging in companion animals" JOURNAL OF ANTI-AGING MEDICINE, vol. 1, no. 2, 1998, pages 117-123, XP000924833 page 117, abstract page 121, column 1, paragraph 3 ---	1-10
A	R.J. KEARNS ET AL.: "Effect of age, breed and dietary omega-6 (n-6) : omega-3 (n-3) fatty acid ratio on immune function, eicosanoid production, and lipid peroxidation in young and aged dogs" VETERINARY IMMUNOLOGY AND IMMUNOPATHOLOGY, vol. 69, no. 2-4, 1999, pages 165-183, XP000924834 AMSTERDAM, NL ISSN: 0165-2427 page 167; table 1 ---	1-10
P,A	WO 99 51108 A (IAMS COMPANY) 14 October 1999 (1999-10-14) claims 1,7 -----	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. l. Application No

PCT/US 00/05586

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9951108 A	14-10-1999	US 5932258 A	03-08-1999
		AU 3378999 A	25-10-1999
<hr/>			